

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

FIG.1





P = polar group N = nonpolar group




FIG. 2

FIG. 3







(XV)



(XVI)




(XVII)




(IVg)

FIG. 4

FIG. 5




IV



m

IVc



IVe

P	NP ¹	R
O-CH ₂ -CH ₂ -NMMe ₂	O-CH ₂ -CH ₂	O-CH ₂ -CH ₂ -CH ₃
O-CH ₂ -CH ₂ -(2-pyridyl)	O-CH ₂ -CH ₂ -NMMe ₂	O-CH ₂ -CH ₂ -CH ₂ -CH ₃
O-CH ₂ -CH ₂ -N(CH ₂ -CH ₂ -NMMe ₂) ₂	O-CH ₂ -CH ₂	O-CH ₂ -CH ₂ -CH ₂ -CH ₃
O-CH ₂ -CH ₂ -(2-imidazolyl)	O-CH ₂ -CH ₂	O-CH ₂ -CH ₂
O-CH ₂ -CH ₂ -NH-Q-NH-NH ₂	O-CH ₂ -CH ₂ -NMMe ₂	O-CH ₂ -CH ₂ -CH ₃
O-CH ₂ -CH ₂ -N(CH ₂ -CH ₂) ₂ -NH	O(CH ₂) ₂ -Me	O-CH ₂ -CH ₂ -CH ₂ -CH ₃

FIG. 6





FIG. 7



P	NP'
$\text{O}-\text{CH}_2-\text{CH}_2-\text{NMMe}_2$	$\text{O}-\text{CH}_2-\text{CH}_3$
$\text{O}-\text{CH}_2-\text{CH}_2-(2\text{-glycidyl})$	$\text{O}-\text{CH}_2-\text{CHMeMe}_2$
$\text{O}-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_2-\text{CH}_2-\text{NMMe}_2)_2$	$\text{O}-\text{CHMeMe}_2$
$\text{O}-\text{CH}_2-\text{CH}_2-(2\text{-imidazolyl})$	OCHMe_3
$\text{O}-\text{CH}_2-\text{CH}_2-\text{NEt}-\text{C}(\text{NEtMe})_2$	$\text{O}-\text{CH}_2-\text{CHMeMe}_2$
$\text{O}-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_2-\text{CH}_2)_2\text{NEt}$	$\text{O}(\text{CH}_2)_2\text{NMe}$






FIG. 8

			Antimicrobial Activity MIC (μg/mL) ¹			Hemolytic Activity HC ₅₀ (μg/mL)
R	X	n	<i>E. c.</i>	<i>K. p.</i>	<i>B. s.</i>	
CH ₂ CH(CH ₃) ₂	(CH ₂) ₂ NHC(=NH)NH ₂	4	20	50	6	200
		5	20	25	6	200
	(CH ₂) ₃ NH ₂	4	12	50	6	200
		5	12	50	6	200
	(CH ₂) ₃ NHC(=NH)NH ₂	4	12	50	12	35
		5	12	50	12	8
	(CH ₂) ₂ NH ₂	2	>60	500	8	>200
		3	>500	>500	37	>200
		4	~30	63	8	>200
		5	100	500	100	
CH(CH ₃) ₂	(CH ₂) ₂ NH ₂	4	100	500	100	
		5	100	500	100	
CH(CH ₃)CH ₂ CH ₃	(CH ₂) ₂ NH ₂	4	500	500	20	
		5	100	500	20	
C ₆ H ₅	(CH ₂) ₂ NH ₂	4	500	500	100	
		5	500	>500	100	
<i>n</i> -C ₄ H ₉	(CH ₂) ₂ NH ₂	4	500	500	500	
		5	100	500	100	
(CH ₂) ₃ NHC(=NH)NH ₂	Me	4	>500	500	500	
		5	500	500	500	
(CH ₂) ₃ NHC(=NH)NH ₂	<i>iso</i> -pentyl	4	100	100	6	4
		5	100	100	12	4
(CH ₂) ₄ NH ₂		2	>500	>500	25	
		4	63	63	<5	

¹ *E.c.* *Escherichia coli* D31; *K.p.* *Klebsiella pneumoniae* 10; *B.s.* *Bacillus subtilis*

FIG. 9

FIG. 10

		Antimicrobial Activity MIC (μg/mL)					
-x-y-	R¹	n	M _n ¹	K91 ²	D31 ³	K.p. ⁴	B.s. ⁵
-CONH-	H	2	756 ⁸	<18	19	66	12
	H	3	1125 ⁸	<19		19	20
	H	10	6000 ⁹	<25	12-50	31-50	1
	H	>30	20,000	>200			5
-NHCONH-	C(=NH)NEt₂		6000		25	100	12.5
	H	2	745 ⁸	<12.5			15
	H	10	6000		<50		5
	H	>30	20000		100		20

¹ *E. coli* K91 (M9 medium)
² *E. coli* K91 (LB medium)
³ *E. coli* D31 (MH medium)
⁴ *Klebsiella pneumoniae* 10 (MH medium)
⁵ *Bacteria subtilis* (LB medium)
⁶ hemolytic activity-erythrocytes HC₅₀ (μg/mL)
⁷ The average chain length is determined by the Flory equation and the polymer size is confirmed by gel chromatography with Waters styryl-gel columns were connected in series to give a MW range from 1,000,000 to 300. The peak was eluted with THF and the peak center at maximum height using a size exclusion column. Average polydispersity for these condensation polymers is ~2.5.
⁸ Homogenous compound prepared by solid phase synthesis.
⁹ *Pseudomonas aeruginosa* 10 IC₅₀ 31-62; *Salmonella typhimurium*, S5 IC₅₀ <3.75; *Enterococcus faecium* IC₅₀ 15-25 (μg/mL).

Time Course of Bacterial Growth





FIG. 11



MIC = 25 µg/ml polyurea

MIC = 5 µg/ml magainin




FIG. 12




FIG. 13























FIG. 14

FIG. 15 **Compound Structures**



Y 093 Z




FIG. 16 **Pmx10073**

FIG. 17 Antagonism of the LMWH Effects on Clotting Time in Whole Blood by Pmx10073.





Table 1. Antibacterial activity and selectivity.




Compound	R ₁	MIC (μg/mL)		HC50 (μg/mL)	Selectivity (HC50/MIC)		Relative Hydrophobicity LogK _{ow}
		<i>E. coli</i>	<i>S. aureus</i>		<i>E. coli</i>	<i>S. aureus</i>	
1	H	12.5	50	12	0.96	0.24	3.51
2		6.25	12	40	6.4	3.3	3.12
3		6.25	6.25	9	1.4	1.4	3.74
4		6.25	6.25	7	1.1	1.1	3.86
5		25	50	790	32	16	1.45
6		25	100	1230 ^a	49	12	2.99
7		50		370	7.4		0.33
8		6.25	12.5	715	110	57	-1.71
MSL-78		12.5		120	9.6		

^aHC50 was obtained from extrapolating the fitted curve to 50 % lysis

FIG. 18

Table 2. Antibacterial activity and selectivity (continued).



Compound	R ₁	R ₂	MIC (μg/mL)		HC50 (μg/mL)	Selectivity (HC50/MIC)		Relative Hydrophobicity Log K _{ow}
			E.coli	S. aureus		E.coli	S. aureus	
1	H	H	12.5	50	12	0.96	0.24	3.51
9	H		25	25	110 nd	4.4	4.4	2.61
10			50	200	400	8.0	2.0	1.53
3		H	6.25	6.25	9	1.4	1.4	3.74
11			12.5	12.5	61	4.9	4.9	2.84
8		H	6.25	12.5	715	110	57	-1.71
12			12.5	12.5	>800	>64	>64	-2.61
MSI-78			12.5		120	9.6		

nd HC50 was obtained from extrapolating the fitted curve to 50 % lysis

FIG. 19




Figure 2. Amphiphilic oligomer 8 induces vesicle leakage.

FIG. 20